

Evaluation of the Relationship between the Level of DHEA-S and Sex Hormones in Some Infertile Iraqi Men

Khaled S. Abd¹ and Adnan F. AL-Azzawie²

¹Salah Al-Din Health Department, Ministry of Health, IRAQ.

²College of Science, University of Tikrit, IRAQ.

¹Corresponding Author: dr_kh683@yahoo.com

ABSTRACT

This study aims to evaluate serum levels of the Dehydroepiandrosterone sulfate (DHEA-S) and its relationship with some sex hormones such as Follicle stimulating hormone (FSH), Luteinizing hormone (LH), prolactin and testosterone hormone and the infertility type in some infertile Iraqi men.

Blood and seminal fluid samples from (60) idiopathic male infertile and (60) healthful individuals as a control group aged (18 to 60 year) were collected from private clinics. Serum hormones (DHEA-S, FSH, LH, prolactin and testosterone) were measurement using Enzyme Linked Immunosorbent Assay (ELISA).

The levels of DHEA-S, FSH, LH and prolactin are significantly higher ($P \leq 0.01$) expect of the testosterone level was significantly decreased ($P \leq 0.01$) in the infertile men as compared with control group. High significant differences ($P \leq 0.01$) were recorded when comparing the hormonal levels (DHEA-S, FSH, LH, prolactin and testosterone) according to the age groups. There are significant differences ($P \leq 0.01$) in levels of (DHEA-S, FSH, LH, prolactin and testosterone) in the infertility period. Smoker infertile men have high levels ($P \leq 0.05$) in the DHEA-S and LH while have low levels in the prolactin hormones compared with control. Patients with family history have shown significant differences ($P \leq 0.05$) in the levels of DHEA-S, FSH, LH and prolactin.

In conclusion, this study revealed significantly increase in the DHEA-S levels in the infertile men and negative correlation between DHEA-S and FSH. Therefore, DHEA-S has important role in the diagnosis and follow up of the male infertility.

Keywords- Male Infertility, DHEA-S, Sex hormones

I. INTRODUCTION

Infertility is the inability to become pregnant after one year of normal, unprotected sexual intercourse with the same partner [1]. Or it is a condition in which males are unable to achieve pregnancy after one year of normal unprotected intercourse, often with a sperm count of less than 20 million/ml [2]. Infertility primarily affects the reproductive system of males and females at equal rates [3].

Dehydroepiandrosterone (DHEA-S) is the most abundant steroid hormone in the human body, and it is called anti-aging hormone or youth hormone because it

delays the aging process [4]. It is a natural steroid hormone consisting of 19 carbon atoms that is produced by body tissues. It is secreted mainly from the zona reticularis of the adrenal cortex. It is also produced to a lesser extent by the brain and gonads, as it is derived from cholesterol as a hormone-generating substance [5].

Natural steroid hormones are derived from cholesterol and can be categorized according to their different physiological roles through activation of their specific intracellular receptors into five classes: glucocorticoids, mineralocorticoids, androgens, estrogens, and progestogens [6].

Follicle-stimulating hormone (FSH) is a member of the glycoprotein family that has a central and essential role in reproduction. (FSH) is a hormone released by the anterior pituitary gland by stimulation of gonadotrophin secretion and possibly other factors. It is released in a pulsatile manner and is partially regulated by glycoproteins, including activin and inhibin. FSH reflects the status of spermatogenesis as a result of feedback between the testis, hypothalamus and pituitary gland [7].

LH is a glycoprotein that regulates testosterone synthesis by Leydig cells. The levels of these hormones are under the control of negative feedback from gonada [8]. In contrast, serum FSH has a longer shelf life, and fluctuations in serum levels are less pronounced. If an abnormal LH value is obtained in one sample, three samples of serum can be collected within a 20-minute interval between one sample and another [9].

Testosterone is responsible for normal growth, development of male reproductive organs, and maintenance of secondary sexual characteristics, as high levels within the testicles are a prerequisite for sperm production and development, improving their motility and improving epididymal function, and there is some controversy about the required relative levels [10].

The hormone prolactin, secreted by the anterior pituitary gland, has a detrimental effect on male fertility when it exceeds the physiological level [11]. It has an important role in the formation of sperm in males, prolactin may have a physiological role in regulating testosterone [12], and hyperprolactinemia leads to infertility in about 11% of males [13]. Where Prolactin controls both LH and FSH production through the regulation of GnRH, as an elevated level of prolactin

will decrease GnRH secretion by slowing down the frequency of GnRH pulsations and thus reducing FSH and LH pulsatility [14].

This study aims to evaluate serum levels of the Dehydroepiandrosterone sulfate (DHEA-S) and its relationship with some sex hormones such as Follicle stimulating hormone (FSH), Luteinizing hormone (LH), prolactin and testosterone hormone and the infertility type in some infertile Iraqi men.

II. MATERIAL AND METHODS

The study included 90 men and included (60) infertile men, and (30) healthy men (control group) at Al-Sharqat General Hospital in Salah El-Din Governorate for the period from October 2020 to the end of December 2020. The samples were divided into age groups. The ages of infertile men ranged from 18 to 60 years and were divided into three groups (less than 20-30 years old, 31-40 years old and over 40 years old). Collected (5) ml was withdrawn from the carpal vein, then blood samples were placed in test tubes free of anticoagulant and left for about a quarter of an hour at room temperature Then it was placed in a centrifuge, G 151 for 10 minutes to obtain the serum, then kept the serum in the freezer in special test tubes at a temperature of -20°C then the serum was frozen at -20°C until the (DHEA-S, FSH, LH, Testosterone and Prolactin) measurement by (ELISA standard curve). The concentrations were quantitatively determined in sera of patients and healthy control subjects by means of ELISA

(Enzyme Linked Immunosorbent Assay) using ready kits manufactured by Bioassay company (China).

Statistical analysis:

Statistical analyses are done using SPSS version 20 computer software. The mean, standard deviation (SD) and of p-value of hormones parameters are calculated using student's t-test and ANOVA test (which considered significant when $p < 0.05$ and highly significant when $p < 0.01$) for the patients and healthy group [15].

III. RESULTS

The effect of infertility on the concentration of hormones:

The results showed that the level of DHEA-S hormone was significantly increased at the probability level of $P \leq 0.01$ among infertile patients (867.9376 ± 21.290) compared with the level of the hormone in normal subjects (763.524 ± 25.726). The results of the current study indicate a decrease in the level of Testosterone hormone 2.5658 ± 0.331 in sterile males when compared with the control group 3.960 ± 0.274 . While it was found that there was a significant increase ($P \leq 0.01$) in the level of hormones FSH 14.620 ± 4.143 (mIu/ml), LH of 8.869 ± 2.646 (mIu/ml) and Prolactin 14.872 ± 2.834 (ng/ml) when compared with the control groups 4.565 ± 2.325 , 4.632 ± 1.110 , 9.303 ± 2.429 , respectively, and as shown in Table (1).

Table 1: Shows comparison between hormonal levels of the infertile and fertile males.

Groups Hormones	Fertility (No. 60)	Infertility (No. 60)	P value
	mean±SD	mean±SD	
DHEA-S	763.524 ± 25.726	867.9376 ± 21.290	≤ 0.01
Testosterone (ng/ml)	3.960 ± 0.274	2.5658 ± 0.331	≤ 0.01
FSH (mIu/ml)	4.565 ± 2.325	14.620 ± 4.143	≤ 0.01
LH (mIu/ml)	4.632 ± 1.110	8.869 ± 2.646	≤ 0.01
Prolactin (ng/ml)	9.303 ± 2.429	14.872 ± 2.834	≤ 0.01

Hormonal levels of infertile males by age:

The results of the current study showed a significant increase ($P \leq 0.01$) in the level of DHEA-S hormone 906.05 ± 12.407 for the age group 31-40 years compared with the hormone level 848.03 ± 9.951 for the age group >30-20 years, and the hormone level 794.89 ± 12.239 for the age group ≥ 41 years. The significant differences ($P \leq 0.01$) in the level of testosterone ng/ml were 2.788 ± 0.356 , 2.71 ± 0.461 , 2.075 ± 0.202 for age groups <30-20 years, 31-40 years

and age group ≥ 41 years, respectively. And statistically significant differences ($P \leq 0.01$) in FSH levels (mIu/ml) 15.490 ± 4.045 , 11.758 ± 4.593 , 18.061 ± 3.495 , respectively, for the same age groups mentioned above. And significant differences ($P \leq 0.01$) in LH levels (mIu/ml) were 8.776 ± 2.302 , 7.648 ± 2.600 , and 10.887 ± 1.375 , respectively. As well as significant differences ($P \leq 0.01$) in the level of prolactin hormone (ng/ml) 17.876 ± 3.338 , 15.427 ± 3.614 , 10.806 ± 2 , respectively. and as shown in Table (2).

Table 2: Shows hormonal and enzyme levels of the infertile males according to age

Age group Hormones	20-30>years (19) sample	31-40 years (25) samples	≥41years (16) sample	P value
	mean±SD	mean±SD	mean±SD	
DHEA-S	848.03±9.951	906.05±12.407	848.03±9.951	≤0.01
Testosterone(ng/ml)	2.788±0.356	2.71 ± 0.461	2.075±0.202	≤0.01
FSH (mIu/ml)	15.490±4.045	11.758±4.593	18.061±3.495	≤0.01
LH (mIu/ml)	8.776±2.302	7.648±2.600	10.887±1.375	≤0.01
Prolactin (ng/ml)	17.876±3.338	15.427±3.614	10.806±2.49	≤0.01

Hormonal levels of infertile males according to the infertility period:

The results of the study showed that the differences were not significant when comparing the level of DHEA-S between age groups, age group ≤10, age group 11-20 and age group ≥21, (850.22±22.678, 804.794±20.459, 945.77±12.761), respectively. Significant differences (P≤ 0.01) in hormone levels of Testosterone (ng/ml) 2.709±0.573, 2.536±0.607, 1.896±0.437. and FSH (mIu/ml) 14.216±2.177,

13.07,5±1.937, 20.455±2.577. and Prolactin (ng/ml) 16.794±1.797, 12.664±1.495, 10.351±0.981. respectively for age groups ≤10 years old, age group 11-20 years and age group ≥21 years. Also, non-significant differences were recorded in the level of LH hormone (mIu/ml) 8.632 ± 1.445, 9.375 ± 1.818, 8.86 ± 1.317, respectively for age groups ≤10 years old, age group 11-20 years and age group ≥21 years. and as shown in Table (3).

Table 3: Shows the hormonal levels of infertile males according to the infertility period

Age group Hormones	Age group ≤10 years (36) sample	Age group 11-20 years (17) sample	Age group ≥21 years (7) samples	P value
	mean±SD	mean±SD	mean±SD	
DHEA-S	945.77±12.76	804.794±20.459	850.22±22.678	≤0.02*
Testosterone(ng/ml)	2.709±0.573	2.536±0.607	1.896±0.437	≤ 0.01 **
FSH (mIu/ml)	14.216±2.177	13.075±1.937	20.455±2.577	0.039*
LH (mIu/ml)	8.632±1.445	9.375±1.818	8.86±1.317	0.04*
Prolactin (ng/ml)	16.794±1.797	12.664±1.495	10.351±0.981	≤0.01

Hormonal levels of infertile males according to smoking:

There were no significant differences (P≤ 0.01) when comparing the level of DHEA-S in smokers (939.446 ± 23.3691) and non-smokers (844.101 ± 48.315). There were significant differences (P≤ 0.01) in the level of LH hormone (mIu/ml) when comparing the level of LH 11.368 ± 2,144 in smokers with the level of LH 8.036 ±2.137 in non-smokers. Also, the differences were significant (P≤ 0.01) in the level of the hormone ng/ml Prolactin when comparing the level of the

hormone 13.308 ± 1.841 in the smokers with the level of the hormone 15.394 ± 2.215 in the non-smokers. The results of the current study did not record significant differences between the level of the hormone Testosterone (ng/ml) when comparing the level of the hormone 2.635 ± 0.559 in smokers with a level of 2.542 ± 0.532 in non-smokers. The same applies to FSH hormone (mIu/ml), as it was found that there were no significant differences in the level of the hormone when comparing between smokers 14,741 ± 2,477 and non-smokers 14.580 ± 2,399. and as shown in Table (4).

Table 4: Shows the hormonal levels of infertile males according to smoking

Groups Hormone	Smokers (No.15)	Non-smokers (No.45)	P value
	mean±SD	mean±SD	
DHEA-S	939.446±23.3691	844.101±48.315	≤0.028
Testosterone (ng/ml)	2.635±0.559	2.542±0.532	0.649

FSH (mIU/ml)	14.741±2.477	14.580±2.399	0.960
LH (mIU/ml)	11.368±2.144	8.036±2.137	0.017*
Prolactin (ng/ml)	13.308±1.841	15.394±2.215	0.029*

Hormonal levels of infertile males by family history:

The results showed that there was a significant difference ($P \leq 0.05$) when comparing the level of DHEA-S 817.34 ± 50.11 in people with a family history with DHEA-S level 874.05 ± 76.05 in people without a family history. The results also showed a significant difference ($P \leq 0.05$) in the level of Testosterone (ng/ml) when comparing the level of the hormone 2.357 ± 0.376 in people with a family history of infertility with the level of the same hormone 2.607 ± 0.323 in people without a family history of infertility. For the same level of significance mentioned above, it was found that there was a significant difference in the level of the hormone Prolactin (ng/ml) when comparing its level of $12.18 \pm$

2.851 among people with a family history of infertility with the level of the same hormone 15.411 ± 3.767 in normal people with no family history of infection. The differences were significant ($P \leq 0.01$) in the level of FSH hormone (mIU/ml) when comparing its level 19.825 ± 2.543 among people with a genetic family history of infertility compared with the level of FSH 13.58 ± 2.073 in normal people without family problems of infertility, as well as for the LH hormone (mIU/ml), it was found that there was a significant difference ($P \leq 0.01$) in the level of this hormone when comparing its level 7.414 ± 1.993 among people with a family history of infertility with the level of the hormone 9.160 ± 1.714 in healthy people and not family history. and as shown in Table (5).

Table 5: Shows the hormonal levels of infertile males by family history

Group Hormones	People with a family history (10) samples	People who do not have a family history (40) samples	P value
	mean±SD	mean±SD	
DHEA-S	817.34±50.11	874.05±76.05	0.02 *
Testosterone(ng/ml)	2.357±0.376	2.607±0.323	0.09
FSH (mIU/ml)	19.825±2.543	13.58±2.073	≤ 0.01 **
LH (mIU/ml)	7.414±1.993	9.160±1.714	0.02 *
Prolactin (ng/ml)	12.18±2.851	15.411±3.767	0.019 *

Correlation between the concentration of DHEA-S with other hormones:

The results of the current study showed a weak positive correlation between DHEA-S, LH ($R=0.075$)

and testosterone ($R=0.006$), while the R correlation of DHEA-S was with FSH ($R=-0.02$) and Prolactin. ($R= -0.082$) Weak negative. and as shown in Table (6).

Table 6: Shows the correlation between the concentration of DHEA-S with other hormones

DHEA-S (ng/ml)	Hormones	Value R
	FSH (mIU/ml)	-0.02
	LH (mIU/ml)	0.075
	Prolactin (ng/ml)	-0.082
	Testosterone (ng/ml)	0.006

IV. DISCUSSION

The effect of infertility on the concentration of hormones:

The results of the current study are in agreement with the study conducted by [16], and the result of the current study is in agreement with the study conducted by [17]. in Iran. The study showed that there was a significant increase in the level of FSH hormone

in infertile men, while the level of testosterone and LH hormone was lower when compared with the control groups. Which showed an increase in FSH hormone with a decrease in Testosterone, while there was no effect on LH. In general, increased FSH levels are a good indicator of germinal epithelial cell destruction and are usually associated with azoospermia or oligospermia. Low testosterone levels are signs of hypogonadism, hypothalamus or pituitary gland [16].

Hyperprolactinemia reduces male fertility by suppressing the action of testosterone, as hyperprolactinemia is well known to lower libido and lead to reduced sperm production, on the other hand, hyperprolactinemia may be caused by a number of pathological factors, such as, disorders of Hypothalamus, pituitary gland tumor, hypothyroidism, and hypogonadism and this may mean some disturbances in the stages of spermatogenesis in infertile patients [18]. The independent secretion of DHEA-S from the adrenal gland and its metabolism in peripheral tissues may explain this finding [19]. The results of the current study in the level of DHEA-S and testosterone hormone and according to the type of infertility agreed with the study [20]. and the study that he conducted [21]. Since testosterone is important in regulating germ cell growth, it becomes necessary to measure the level of testosterone to monitor male infertility [22]. Sperm formation requires the presence of gonadotropins and testosterone [23]. Spermatogenesis is associated with higher levels of intracellular testosterone, and our results were in agreement with that of [24], thus not only explaining impaired spermatogenesis but also reduced function of Leydig cells and causes of pretesticular azoospermia were also suspected [24]. And it was shown by a previous study that was conducted that there is no direct correlation between the level of DHEA-S in the serum and the number of sperms [21], the independent secretion of DHEA-S from the adrenal gland and its metabolism in the peripheral tissues may explain this result [19].

Hormonal levels of infertile males by age:

The influence of age on infertility varies and interacts with many genetic, social, psychological and health factors related to the patient's history [25]. And it did not find an association or effect on the age factor with infertility, as this was not statistically significant, while the results of [26]. It found that males had a higher rate of infertility between 20-29 years (46%), followed by 30-39 years (40%). While [27]. specifically indicated in his study that the incidence of infertility at the age of less than 30 years is higher in men, followed by 30-34 years and decreases with age. The results of a study also found that the vast majority of sterile patients ranged in age from 20-29 years [28]. Their results were attributed to early marriage, semen quality, and ejaculation process, which gradually decrease with age. It begins to decline after the age of 35 [29]. As well as the presence of many other genetic disorders that have an effect on the reproductive process, as is the case in the X chromosome [30]. DHEA-S can turn into an androgen or estrogen, and not all of its vital functions have been determined so far. The concentration of DHEAS in the blood reaches its peak around the age of 25 years and then declines steadily during the following years, due to its half-life. With its long duration and high concentration in the blood, DHEAS levels remain the same 24 hours a day. This makes DHEAS an important

diagnostic tool for both scientific research and clinical diagnosis. Furthermore, concentrations of DHEAS can change by many factors, such as production Internal medicine, hormonal supplements, many types of medications, and many types of disease conditions [31].

Hormonal levels of infertile males according to the period of infertility:

The effect of post-marital infertility differed on hormonal behavior. Although no studies have been conducted on the effect of this factor on concentrations of specific hormones, the effect of age can depend mainly on the fact that testosterone tends to decrease with increasing infertility as well as an increase in the level of FSH. These findings can also be associated with poor mental status and emotional distress in patients.

Hormonal levels of infertile males according to smoking:

The results of the current study agreed with the results of the study conducted by [32], and differed with the results of the study conducted by [33]. Many other studies have shown that the relationship of smoking and male infertility is uncertain, ie. The issue of smoking and its effect on infertility did not have a clear effect and the results of studies varied from one study to another [34], and studies showed that there are no reliable effects among male smokers on semen parameters. Nicotine may modulate the pituitary axis by enhancing the production of cortisol, growth hormone, oxytocin and vasopressin, which in turn inhibit prolactin and luteinizing hormone [35]. In a study looking at the effectiveness of tobacco smoking on hormone levels [36], average levels of estradiol were observed to be higher, and the average levels of prolactin, follicle-stimulating hormone (FSH) and LH levels were lower in smokers compared to non-smokers, while the average levels of FSH were not. Testosterone and dehydroepiandrosterone are different. Many components of tobacco smoke have effects on testosterone. Nicotine has been reported to impair the male reproductive hormone system by causing Leydig apoptosis and inhibiting androgen synthesis [37]. Toxins, including lead in tobacco, appear to directly impair spermatogenesis itself as well as sperm function through reproductive axis impairment or testicular degeneration. Also, semen parameters (total number, motility, and morphology) are decreased in infertile smokers compared to infertile non-smokers. Smoking caused a significant decrease in sperm count and morphology. However, it had no significant effect on movement in the study subjects [38].

Hormonal levels of infertile males by family history:

It included several subgroups of infertile subjects which in all cases indicated no family history of infertility. In their survey, this included the chromosomal assessment of the members of the infertility group studied [39]. While some studies have indicated that families with a medical history of infertility and a family history of infertility called PRM1 homozygous PRM1

genetic mutations have a 40% higher risk of infertility than those with no family history of infertility [40].

Correlation between the concentration of DHEA-S with other hormones:

The results of the current study agreed with the results of the study conducted by [41], that an increase in the concentration of DHEA-S hormone is associated with a decrease in the concentration of testosterone in the blood [42]. In a study conducted, it was found that there is an elevated chromatin damage in men with low or low androgen status. DHEA and its sulfate ester (DHEA-S) are the two most abundant steroid hormones in humans [43].

In conclusion, this study revealed significantly increase in the DHEA-S levels in the infertile men and negative correlation between DHEA-S and FSH. Therefore, DHEA-S has important role in the diagnosis and follow up of the male infertility.

REFERENCES

- [1] Turek, PJ (2008) Male infertility. In: Tanagho EA, Mc Aninch JW (Eds.), (17th edn) Smith's general urology, Mc Graw-Hill Companies, New York, USA, pp. 684-717.
- [2] Wong WY, Thomas CM, Merkus JM, Zielhuis GA, Steequeers-Theunissen RP (2000) Male factor sub fertility: Possible causes and the impact of nutritional factors. *Fertil Steril* 73(3): 435-442.
- [3] Sohrabvand F, Jafarabadi M (2005) Knowledge and attitudes of infertile couples about assisted reproductive technology. *Int J Reprod Biomed* 3(2): 90-94.
- [4] Abraham, P.A.; Kazman, J.B.; Zeno, S.A.; Poth, M. and Deuster, P.A. (2013). Age-related decline in salivary dehydroepiandrosterone sulfate and associated health risks among African Americans. *Ethn Dis.* 23(2):149-54.
- [5] Ross, M and Pawlina, W. (2011). *Histology: A Text and Atlas* (6th ed.). Lippincott Williams & Wilkins. pp.708,780.
- [6] Franchimont, P. 1983. Regulation of gonadal androgen secretion. *Horm Res* 18:7-17.
- [7] Jarow, J., Sigman, M., and Kolettis, P. N. (2010). AUA-Clinical guidelines: the optimal evaluation of the infertile male: best practice statement. AUA-Home. American Urological Association Education and Research, Inc.
- [8] Jarow, JP. (2003). Endocrine causes of male infertility. *Urol Clin North Am.* 30: 83-90.
- [9] Goldzieher JW, Dozier TS and Smith KD. (1976). Improving the diagnostic reliability of rapidly fluctuating plasma hormone levels by optimized multiple-sampling techniques. *J Clin Endocrinol Metab*; 43:824.
- [10] Hameed S, Jayasena CN, Dhillon WS. (2011). Kisspeptin and fertility. *J Endocrinol.* 208:97-105.
- [11] Singh R., H Alaa and Agarwal A. (2011). Thyroid Hormones in Male Reproduction and Fertility *Rajender. The Open Reproductive Science Journal*, 3, 98-104.
- [12] Sigman, M. and Jarow, JP. (2007). "Current Management of Male Infertility, An Issue of Urologic, E-Book". Saunders Elsevier. ed.10th; p. 609-53, 6.
- [13] Masud, S; Mehboob, F; and Bappi MU; (2007). "Severe hyperprolactinemia directly depresses the gonadal activity causing infertility". *Esculapio J Services Inst Med Sci.* 2:25-7.
- [14] Maria, M.; (2016). The Role of Prolactin in Men. *Endocrinol Metab Syndr.* 5: 222.
- [15] Al-Rawi, K, M. (2000). An introduction to statistics. Second Edition. House of Books for Printing and Publishing - University of Mosul.
- [16] Zalata, A. A. et al. (2008) 'Follicle-stimulating hormone receptor polymorphism and seminal anti-Müllerian hormone in fertile and infertile men', *Andrologia.* Wiley Online Library, 40(6), pp. 392-397.
- [17] Safarinejad, M. R., Shafiei, N. and Safarinejad, S. (2011) 'Evaluating the role of the FSH receptor gene Thr307-Ala and Asn680-Ser polymorphisms in male infertility and their association with semen quality and reproductive hormones', *Bju International.* Wiley Online Library, 108(2b), pp. E117-E125.
- [18] Al-Nahi, A. S. (2015) 'Evaluation of FSH, LH, Testosterone, Prolactin, TSH and T4 hormones levels in different subgroups of infertile males.', *Al-Kufa University Journal for Biology.* University of Kufa, 7(3), pp. 47-54.
- [19] Phillips GB, Relationship between serum dehydroepiandrosterone sulfate, androstenedione, and sex hormones in men and women. *European journal of endocrinology.* 1996; 134 (2): 201-6.
- [20] Al-Siddeig, M. and Ismail, A. (2016). Serum Testosterone and Dehydroepiandrosterone-Sulfate (DHEA-S) Level among Oligospermia and Azoospermia Patients. Volume 15, Issue 4; PP 62-65.
- [21] Garcia Diez, LC, Gonzalez Buitrago JM, Corrales JJ, Battaner E, Mirales JM. Hormone level in serum and seminal plasma of men with different type of azoospermia. *J Reprod Fertile* 1983; 67: 209-14.
- [22] Ramesh, Babu S., Sadhnani M.D., Swarna M., Padmavathi P. and Reddy P.P. (2004). Evaluation of FSH, LH and testosterone levels in different subgroups of infertile males. *Indian Journal of Clinical Biochemistry*, 19 (1) 45-49.
- [23] Lalitha c, Sayee R, Jeyanthi K, Shubha R. Hormonal factors associated with hypogonadism and infertility in males – chromosomal abnormality. *IOSR Journal of Dental and Medical Sciences.* 2013; 10 (1): 71-75.
- [24] Masanori Yamamoto, Satoshi Katsuno, Keisuke Yokoi, Hatsuki Hiei, and Koji Miyake, Serum inhibin levels in normal men and men with idiopathic infertility. *Nagoya J. Med. Sci.* 1995; 58: 35-39.
- [25] Mossa, H. A. L. (2010) 'The Effect of Seasonal, Age, Smoking and Occupational exposure status on semen parameters in Iraqi Subjects', *Al-Mustansiriyah Journal of Science.* Al-Mustansiriyah University, 21(7), pp. 12-18.

- [26] Khadhim, M. M., Al-quizwini, T. S. and Hussian, T. A. (2015) 'The Role of Protamine 1 Gene Polymorphism and Anti FSH Antibody in Infertile Couples', *Al-Qadisiyah Medical Journal*, 11(20), pp. 85–91.
- [27] Al-Haija, R. W. M. A. (2011) 'Main causes of infertility among men treated at Razan Centers in West Bank: Retrospective study'.
- [28] Moghadam, M. H. B. et al. (2011) 'Evaluation of the general health of the infertile couples', *Iranian journal of reproductive medicine. Shahid Sadoughi University of Medical Sciences and Health Services*, 9(4), p. 309.
- [29] Omran, H. M., Bakhiet, M. and Dashti, M. G. (2013) 'Evaluation of age effects on semen parameters of infertile males', *Andrology*, 2(106), pp. 250–2167.
- [30] Raman, J. D. and Schlegel, P. N. (2002) 'Aromatase inhibitors for male infertility', *The Journal of urology. Wolters Kluwer Philadelphia, PA*, 167(2 Part 1), pp. 624–629.
- [31] Leowattana, W. (2004). DHEAS as a new diagnostic tool. *Clinica Chimica Acta; International Journal of Clinical Chemistry*, 341, 1–15.
- [32] Boeri, L., Capogrosso, P., Ventimiglia, E., Pederzoli, F., Cazzaniga, W., Chierigo, F., et al. (2019). Heavy cigarette smoking and alcohol consumption are associated with impaired sperm parameters in primary infertile men. *Asian J. Androl.* 21, 478–485. doi: 10.4103/aja.aja_110_18.
- [33] Rehman R, Zahid N, Amjad S, Baig M and Gazzaz ZJ (2019) Relationship Between Smoking Habit and Sperm Parameters Among Patients Attending an Infertility Clinic. *Front. Physiol.* 10:1356.
- [34] Wallach, E. E., Hughes, E. G. and Brennan, B. G. (1996) 'Does cigarette smoking impair natural or assisted fecundity?', *Fertility and sterility. Elsevier*, 66(5), pp. 679–689.
- [35] Weisberg, E. (1985) 'Smoking and reproductive health.', *Clinical reproduction and fertility*, 3(3), p. 175.
- [36] Ochedalski, T. et al. (1994) 'Examining the effects of tobacco smoking on levels of certain hormones in serum of young men', *Ginekologia polska*, 65(2), pp. 87–93.
- [37] Dai, JB, Wang ZX. and Qiao ZD. (2015). The hazardous effects of tobacco smoking on male fertility. *Asian J Androl.* 17(6):954–60.
- [38] Gandhi, J., Hernandez, R. J., Chen, A., Smith, N. L., Sheynkin, Y. R., Joshi, G., et al. (2017). Impaired hypothalamic-pituitary-testicular axis activity, spermatogenesis, and sperm function promote infertility in males with lead poisoning. *Zygote* 25, 103–110.
- [39] Babu, S. R. et al. (2004) 'Evaluation of FSH, LH and testosterone levels in different subgroups of infertile males', *Indian Journal of Clinical Biochemistry. Springer*, 19(1), pp. 45–49.
- [40] Jawad, E. S., Mubark, H. R. and Odah, T. K. (2019) 'The Relationship Between The Genotypes Frequency For The PRM1 Gene And Some Of Risk Factors In Male Infertile Of Thi Qar Province', *Journal of Education for Pure Science. Thi-Qar University*, 9(1), pp. 102–112.
- [41] Ismael, Z.K, M.O. Selman, H.A.L. Mossa and L.A. AL-Anbari (2017). Relationship of FSH, LH, DHEA and Testosterone Levels in Seminal Plasma with Sperm Function Parameters in Infertile Men. *Merit Research Journal of Medicine and Medical Sciences*, 5(12): 627-634.
- [42] Cannarella, R.; Condorelli, R.; Dall'Oglio, F.; Vignera, S.; Laura M. Mongio, L.; Micali, G.; and Calogero, A. (2020). Increased DHEAS and Decreased Total Testosterone Serum Levels in a Subset of Men with Early-Onset Androgenetic Alopecia: Does a Male PCOS-Equivalent Exist?. *International Journal of Endocrinology. Volume 2020. P 8.*
- [43] Charlton, M., Angulo, P., Chalasani, N., Merriman, R., Viker, K., Charatcharoenwitthaya, P., Sanderson, S., Gawrieh, S., Krishnan and A., Lindor, K. Low circulating levels of dehydroepiandrosterone in histologically advanced nonalcoholic fatty liver disease. *Hepatology.* 2008; 47:484–492.